


## Tyrosine Hydroxylase Rabbit mAb (AR1580)


## **Key Features**

Host Species:	Rabbit		
Reactivity:	Human, Mouse, Rat		
Applications:	WB,IHC,IF,IP,ELISA		
lsotype:	IgG,Kappa		
MW:	59kD (Calculated) 59kD (Observed)		
Recommended Dilution Ratios			
IHC:	1:500-2000		
WB:	1:2000-10000		
IF:	1:200-1000		
ELISA:	1:5000-20000		
IP:	1:50-200		
Storage	-15°C to -25°C/1 year (Do n	ot lower than -25°C)	
<b>Basic Information</b>			
Clonality	Monoclonal		
Immunogen Information			
Specificity	Endogenous		
Target Information			
Gene name	ТН ТҮН		
Protein Name	Tyrosine 3-monooxygenase (Tyrosine 3-hydroxylase) (TH), Tyrosine Hydrolase		
	Organism	Gene ID	UniProt ID
	Human	7054	P07101
	Mouse	21823	P24529
	Rat	25085	P04177
Tissue specificity	Mainly expressed in the brain and adrenal glands.		

## Validation Data







Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-Tyrosine Hydroxylase antibody. The HRP conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: Rat brain Lane 2: Mouse brain Predicted band size: 59kDa Observed band size: 59kDa

Human brain was stained with anti-Tyrosine Hydroxylase rabbit antibody

Immunofluorescence analysis of Hela cell.

- TH Monoclonal Antibody(red) was diluted at 1:200(4 ° overnight). GAPDH Monoclonal Antibody(2B8)(green) was diluted at 1:200 (4° overnight).
- 2. Goat Anti Rabbit Alexa Fluor 594 was diluted at 1:1000(room temperature, 50min). Goat Anti Mouse Alexa Fluor 488 was diluted at 1:1000(room temperature, 50min).

For Research Use Only